

KIVILIS, S.S.; KUZNETSOVA, M.I., red.

[Regulations 28-64 for measuring the consumption of fluids, gases and steam by standard diaphragms and nozzles] Pravila 28-64 izmereniia raskhoda zhidkostei, gazov i parov standartnymi diafragmami i soplami. Izd. ofitsial'noe. Moskva, Izd-vo standartov, 1964. 146 p. (MIRA 18:2)

1. Russia (1923- U.S.S.R.) Komitet standartov, ser 1 izmeritel'nykh priborov.

KIVILIS, S.S.

[Regulations 28-64 on the measurement of the consumption of liquids, gases and vapors by standard diaphragms and nozzles] Pravila 28-64 izmereniia raskhoda zhidkostei, gazov i parov standartnymi diafragmami i soplami. Izd. ofitsial'noe. Moskva, Izd-vo Standartov, 1964. 148 p. [Album of diagrams for....] Al'bom grafikov k.... (MIRA 18.5)

1. Russia (1923- U.S.S.R.) Komitet standartov, ser 1 izmeritel'nykh priborov.

KIVILIS, S.S.; GAFANOVICH, M.D.

Measurement of consumption by pressure drop with an automatic
compensation of gas parameter changes. Izv. tekhn. no. 12:40-45
D '64. (MIRA 18:4)

L 2264-66 EWT(4)/EWP(v)/EWP(h)/EWP(h)/EWP(1)
ACCESSION NR: AP8011068

UR/0115/68/000/003/0052/0054

AUTHOR: Kivilis, S. S.; Reshetnikov, V. A.

TITLE: The effect of steady-state flow profile on the error of ultrasonic flow-meters

SOURCE: Izmeritel'naya tekhnika, no. 3, 1968, 52-54

TOPIC TAGS: hydromechanics, hydraulic engineering, ultrasonic flow meter, pipeline flow, ultrasound, flow profile, velocity distribution curve

ABSTRACT: The authors discuss the effect of the flow profile contour on the systematic error of flow-meters, noting that this is one of the most important problems in the measurement of flow rates by means of ultrasound. It is noted that when determining the rate of flow of a liquid passing through a pipeline, it is essential to know the velocity averaged over the flow cross section. These averaged velocities are related by a non-linear function, the analytical expression for which (in the case of a cylindrical pipeline) is presented and analysed in the article. Attention is called to the fact that, while this expression and others similar to it suggested by various authors are based on a logarithmic law for the distribution of velocities in the pipeline, when $x = \text{const}$, this law is merely an approximate description of the actual velocity distribution curves and the constant x , used in the expression of the logarithmic law, changes even within the limits

Card 1/3

L 2264-66

ACCESSION NR: AP5011063

of a single velocity distribution curve for a constant Reynolds number. The authors therefore attempt to determine the effect of the velocity profile in the stream on the accuracy of ultrasonic flow-meter readings by means of a direct integration of experimentally derived velocity distribution curves. Expressions are obtained for the mean velocity of a flow of any configuration and for the mean velocity along the ultrasound propagation path. In this way, an accurate formula is derived describing the relationship between these velocities for a flow of any configuration. This expression is modified for the particular case of a steady-state stream in a cylindrical pipeline of given radius, with the ultrasound propagating in a plane which passes through the axis of the pipe. Steady-state flow velocity distribution curves are given in the article for different Reynolds numbers running from $4 \cdot 10^3$ to $3 \cdot 10^5$. A Chebyshev formula was used as the working formula in the integration of velocity distribution curves, with the basic data taken from the table of Nikuradze (Problemy turbulentnosti. ONTI, M.-L., 1936). The results of this computational work are presented in a separate figure, from which it is clear that for Reynolds numbers up to approximately $20 \cdot 10^3$ there is good agreement between the results obtained using the equations given in the first section of the article and the data obtained by direct integration of the velocity curves. Finally, an empirical formula is offered, expressing the quantity m as a function of the Reynolds number and obtained on the basis of the distribution integration carried out previously. This formula is simple and may be used in practical computations. Orig. art. has: 2 figures and 11 formulas.

Card 2/3

L 2264-66

ACCESSION NR: AP6011008

ASSOCIATION: None

SUBMITTED: 00

ENCL: 00

SUB CODE: ME, IE

NO REF SOV: 008

OTHER: 001

Card

3/80

1-2508-66 EWT(d)/EWT(1)/EWC(k)-E/EWP(v)/EWP(k)/EWP(h)/EWA(h)/EWP(1)
 ACCESSION NR: AP5019201 UR/0115/65/000/006/0053/0057
 531.732 / 3 (083)

AUTHOR: Kivilla, S. S. qm

TITLE: Measuring rate-of-flow of liquids and gases by restriction-type differential manometers 25

SOURCE: Izmeritel'naya tekhnika, no. 6, 1965, 53-57

TOPIC TAGS: manometer, differential manometer 14

ABSTRACT: Fundamentals of the new "Rules 28-64 for measuring rate-of-flow of liquids, gases, and steam by standard orifice plates and nozzles" are discussed. Connected with the International Standard Organization (ISO) provisions, these "Rules" are applicable to measuring the flow of single-phase liquids, gases, and superheated steam by a restriction mounted inside a pipe 50-mm or more in diameter, for a steady-state flow, certain Re numbers and pressure ratios. The Venturi tubes have not been standardized as yet. Saturated

Card 1/2

L 2598-66

ACCESSION NR: AP5019201

steam and pulsating flows are not to be measured with differential manometers. Mass flow, instead of weight flow, is dealt with. Formulas, curves, and tables for designing restrictions are provided in the "Rules"; also, the Section on gas compressibility has been "materially reworked." Nomographs are supplied for determining the optimal nominal pressure drop caused by a restriction. International requirements of restrictions are adopted. Methods for determining the measurement error have been developed. Orig. art. has: 3 figures, 13 formulas, and 1 table.

ASSOCIATION: none

SUBMITTED: 00

ENCL: 00

SUB CODE: IE, ME

NO REF SOV: 008

OTHER: 012

mk
Card 2/2

...meter (approximately 1.5 mm diameter)

...meter

ABSTRACT An Author Certificate has been issued for a flowmeter, see Fig. 1 of the ... based on the measurement of the variation of pressure in the flow. The range of measurement is ... a swirling element in the flow ... the pressure in the flow tube and a tube for measuring the pressure ... periphery. Orig. art. has: 1 figure.

(AC)

PRINTED 15Aug64

ENCL: 01

SWR CODE: ME, IE

NO REF SOV: 000

OTHER: 000

ATD PRESS: 4037

Card 1/2

10095-5

ACCESSION NR: AP5016749




Fig. 1. Flowmeter

1 - Swirler; 2 - core
pressure tube; 3 - peri-
phery pressure tube.

llc
Card 2/2

KIVILIS, S.S.; RESHETNIKOV, V.A.

Hydromechanical error of ultrasonic flowmeters. Izv. tekhn.
no.11:48-50 N '65. (MIRA 18:12)

L 35845-66 EWT(1) JAJ

ACC NR: AP6014523

SOURCE CODE: UR/0115/65/000/011/0048/0050

AUTHOR: Kivilis, S. S.; Reshetnikov, V. A.

59

ORG: none

B

TITLE: Hydromechanical error of ultrasonic flowmeters

SOURCE: Izmeritel'naya tekhnika, no. 11, 1965, 48-50

TOPIC TAGS: fluid mechanics, ultrasonic equipment, flow meter, ~~ERROR~~
CORRECTION

ABSTRACT: Presently known types of ultrasonic flowmeters can be divided into three groups, depending on the velocity of the flow being measured. The article gives a table showing the main characteristics of these types, using the following nomenclature: L_0 is the distance between the radiating and receiving piezo-transformers; $u(L)$ is the distribution of the flow velocity along the propagation path of the ultrasound; v_0 is the flow velocity along the axis of the pipeline; v_m is the maximum flow velocity; r_0 is the radius of the pipeline. Keeping in mind that the correction coefficient

$$K = \frac{v}{v_0} \quad (1)$$

where v is the velocity to be measured; v_0 is the average velocity over

Card 1/2

UDC: 531.732.083

Card 2/2

KIVILO, Evald; RAIG, H., otv. red.

[Elementary course in hygiene for workers of dairy
farms] Sanitaarmiinimumi kursus piimakarjafarmide
töötajatele. Tartu, Vabariiklik sanitaarhariduse
maja, 1962. 61 p. (MIRA 16:12)

KIVILO, M.; PURDE, M.

Use of semiautomatic inhalation anesthesia experiments on animals.
Biul. eksp. biol. i med. 58 no.10:124-125 O '64.

(MIRA 18:12)

1. Estonskiy respublikanskiy onkologicheskiy dispanser (glavnyy vrach A.N. Gavrilov) i Estonskiy institut eksperimental'noy i klinicheskoy meditsiny (dir. - prof. P.A. Bogovskiy) AMN SSSR.
Submitted July 15, 1963.

KIVIL'SHA, Ye.A.

Treatment of otogenous cerebral abscesses. Vest.oto-rin. 17 no.1:
60 Ja-F '55. (MIRA 8:5)

1. Iz otdeleniya bolesney ucha, gorla i nosa bol'nitay in. Lenina,
Kamenets-Podol'skiy.
(BRAIN-ABSCESS)

KIVILSHA, I. E., Cand of Tech Sci -- (diss) "On the Problem of the Use of
Local Cement Made from Chalky Marl," Kaunas, 1959, 31 pp (Kaunas
Polytechnical Institute) (KL, 2-60, 113)

METSIK, R.; TOMBERG, A.; RAYAVEE, E. [Rajavee, E.]; KIVIMAA, Kh. [Kivimaa, H.]

Investigating phenols extracted from semicoking shale tars by sodium carbonate aqueous solutions. Khim. i tekhn.gor.slav. i prod. ikh perer. no.12:181-192 '63. (MIRA 17:2)

SHELOUMOV, V.V.; KIVIMAA, H.M. [Kivimaa, H.]

Centrifugation of heavy shale tars. Khim. i tekhn. gor. slan. i prod.
ikh perer. no.11:220-229 '62. (MIRA 17:3)

SHELOUMOV, V.V.; METSIK, R.E.; KAL'BERG, A.O. [Kalberg, A.];
KIVIMAA, Kh.M. [Kivimaa, H.]

Preparing oil shale tar for distillation. Khim. i tekhn. gor.
slan. i prod. ikh perer. no.10:174-190 '62. (MIRA 17:5)

KHYUSSE, I.Yu.; SHELOUMOV, V.V.; RAYAVEYE, E.L.; METSIK, R.E.; KIVIMAA, Kh.M.
(Kivimaa, H.)

Certain possibilities of increasing water soluble phenol resources.
Khim. i tekhn. gor. slan. i prod. ikh perer. no.11:230-235 '62.
(MIRA 17:3)

KIVIMYAGI, E.A.; YUDOVICH, E.A.

Amazine treatment of patients with schizophrenia suffering from
pulmonary tuberculosis. Zhur.nevr.i psikh. 61 no.2:247-250 '61.
(MIRA 14:6)

1. Ryssanskaya psikhonevrologicheskaya bol'nitsa (glavnyy vrach
V.V.Tsarichenko, vypolnena pod rukovodstvom prof. A.K.Strelyukhina).
(TUBERCULOSIS) (SCHIZOPHRENIA)
(CHLORPROMAZINE)

KIVIN, Abram Maturovich, inzhener; UGAROV, I.P., ishener redaktor;
KANDYKIN, A.Ye., tekhnicheskiy redaktor

[Running trains without stopping to take on water; a collection
of articles] Vozhdenie poezdov bez ostanovok dlia nabora vody;
sbornik statei. Moskva, Gos.transp.shel-dor.isd-vo, 1955. 42 p.
(Railroads--Water-supply) (MLRA 8:10)

YEVDOKIMOV, I.I.; ALEKSEYEV, V.D.; ASHIKHEMIN, A.K.; BAYEV, M.V.; BEGLAR'YAN, P.A.; BYCHKOV, I.A.; VASLOVA, Ye.T.; VYZHEKHOVSKAYA, M.P.; GURETSKIY, S.A.; DEMIDOV, I.M.; YESSIOV, Ye.P.; ZHUKOV, V.D.; ZELINSKIY, M.G.; ZOL'NIKOV, F.T.; ZOLOTOVA, L.I.; KIVIN, A.N.; KOMARNITSKIY, Yu.A.; KONSTANTINOV, A.N.; KUL'CHITSKAYA, A.K.; MAKSIMENKO, I.I.; MELEST'YEV, A.A.; MOROZOV, I.G.; MURZINOV, M.I.; OZEMBLOVSKIY, Ch.S.; OSTRYAKOV, K.I.; PANINA, A.A.; PAVLOVSKIY, V.V.; PERMINOV, A.S.; PERSHIN, B.F.; PRONIN, S.F.; PRIMENNYI, A.I.; POKHOVSKIY, M.I.; RASPOVOMAREV, Ye.A.; SEMIN, I.N.; SKLYAROV, Yu.N.; TIBABSHIN, A.I.; FAREKROV, Ye.D.; YNDOROV, G.P.; SHUL'GIN, Ye.S.; YAKIMOV, I.A.; VERINA, G.P., tekhn.red.

[Labor feats of railway workers; stories about the innovators]
 Trudovye podvigi shelasnodorozhnikov; rasskazy o novatorakh. Moskva,
 Gos.transp.shel-dor.isd-vo, 1959. 267 p. (MIRA 12:9)
 (Railroads) (Socialist competition)

Methods for testing cold-chamber materials. N. A. Zhikharevich and D. I. Kargin. (Zhurnal. Tekhn.-fizi-
chesk. iust. Opyt. i teo. Khim. 32, 112 pp. (1955).
—A series of rational methods was developed for detg. the
size of bodies in acids, and for measuring their d. The
work was based on the Kallman-Boris method suppli-
mented by (a) dissolving the tiny sulfuric crystals by
heating them on a water bath, (b) neutralizing "acid"
grains (grains treated with H_2SO_4) in the presence of methyl
orange, (c) neutralizing "alkali" grains (grains treated
with Na_2CO_3) in the presence of phenolphthalein.

M. V. Kondalov

KIVIN, D. I.

Jun 1947

USSR/Engineering
Metallurgical Plants
Dolomite

"Dolomite Bricks in Metallurgy," Prof G. V. Kukolev, Dr Tech Sci; D.I. Kivin, Engr, All-Union Inst Fireproof Materials, 5 pp

"Stal'" No 6, vol. 7, p. 531

Use of dolomite bricks in important elements of furnaces was unsatisfactory because of their shrinkage and deformation due to high temperatures. From experiments, high-quality, water-resistant dolomite brick developed to replace magnesium and chrome-magnesium bricks. New brick will effect on quantity and quality of steel casting for present Five-Year Plan, since dolomite resources are available at almost all metallurgical processing areas.

PA 58134

Dominance of water-resistant dinitamite both in service. G. V. Kishiyev and D. I. Kishin, *Chemistry* 12, 207-21 (1967), cf. *C.A.B.* 65, 877-8; 67, 1464-5; 68, 1464-5. The new dinitamite polyacrylates are even better than dinitamite-maleic anhydride. It is claimed that their stability is superior to the usual dinitamite-maleic anhydride. Their acid resistance in open-laminate conditions is excellent, owing to the development of a rather plastic structure. No swelling occurs, certainly in parallel direction, and laminates, in which their life is 5 to 8 times that of dinitamite brick. Chem. analysis shows the changes in composition, in d_n , and in strength of the different zones after service. The CaO ratio, $\text{mol.} / (\text{KCN})$ is characteristic for these zones as an indication of their migration, especially of the surface, where the CaO is the highest parts. A relative enrichment of MgO is found with this phenomenon. However, better study of Fe_2O_3 dinitamite and 50% polyacrylolytic composite shows in the hottest zone, a relative decrease in MgO and enrichment in CaO . In H_2O -resistant dinitamite polyacrylates the Fe_2O_3 content must be low to reduce staining, the MgO (as the most refractory oxide) must be high, and the CaO ratio

must be as high as possible, but not near 1.8, in order to avoid a CaO crust. Both open-hearth slags which react with the dolomite or magnesia brick lining affect not only the chem. compn. of the products but also the refractory character, due to the phase equil. in the system $\text{SiO}_2\text{-MgO-CaO}$ in its basic part. The most characteristic secondary phases are $\text{SiCa}_2\text{Si}_2\text{O}_8$, $\text{SiCa}_2\text{Si}_2\text{O}_8$, hematite, magnetite, spinels (with FeO and MnO in cryst. soln.), Ca_2SiO_4 , Ca_2FeO_4 , $\text{SiCa}_2\text{Al}_2\text{O}_8$, $\text{SiCa}_2\text{Al}_2\text{O}_8$, spinels, hematite, and $\text{SiCa}_2\text{FeO}_8$ in the slags. The Si factor is decreased from 0.55-0.58 to 0.57 in dolomite, while in magnesia it is about 0.55 after service. The decrease of $\text{SiCa}_2\text{Si}_2\text{O}_8$ to $\text{SiCa}_2\text{Si}_2\text{O}_8$, and free CaO and a reduction in peroxide content are characteristic of conditions with slag, but the amt. of fused material is still low. This explains the excellent stability of dolomite with or without CaO to 1700°. The $\text{SiCa}_2\text{Si}_2\text{O}_8$ in the original lining also acted as a buffer against the formation of Ca_2SiO_4 during service as a buffer against the formation of Ca_2SiO_4 during service. Ca Mg silicates. The decrease of $\text{SiCa}_2\text{Si}_2\text{O}_8$ below 1350-1400° does not cause disintegration because the original slag-reaction zone protects the free CaO from hydrolysis. W. Hill

SOV/81-59-9-32088

Translation from: Referativnyy zhurnal. Khimiya, 1959, Nr 9, p 358 (USSR)

AUTHORS: Kukolev, O.V., Kivin, D.I., Zelenskaya, A.T., Lur'ye, M.A., Minskiy, Ya.M.

TITLE: Magnesite-Dolomite Highly-Refractory Products

PERIODICAL: Sb. nauchn. tr. Vses. n.-i. in-ta ogneporov, 1958, Nr 2 (49), pp 277 - 296

ABSTRACT: The manufacture of magnesite-dolomite products from clinkers with various content of dolomite (D) and magnesite (M) in the raw material mixture of the clinker has been studied. Sotka M and Karagay D served as raw material; for binding CaO, crystalline quartzite and iron scale were introduced; for the stabilization of β -2CaO · SiO₂ an addition of phosphorite ore was introduced. The composition of the magnesite-dolomite charge was so calculated that a high (~ 1) coefficient of saturation with lime was obtained. Four charges were prepared: I - the ratio of M to D = 1:1; I² - the same with an increased content of scale, II and III with the ratio M to D = 1:2 and 2:1, respectively. Dried briquets from charges I, I² and II were burnt in the rotating furnace

Card 1/2

Magnesite-Dolomite Highly-Refractory Products

SOV/81-59-9-32088

at 1,710 - 1,760°C and from charge III in the periodic furnace at 1,600°C; the burnt briquets were ground and from the powders (the grain composition is cited) products were formed and burnt: from charges I, I^P and II at 1,430°C, from charge III at 1,460°C. A part of the raw bricks were left for hydraulic hardening for obtaining bricks without burning. The bricks from all the charges, in spite of the low burning temperature, have a high density (porosity 8.12 - 14.1%), high mechanical resistance (σ_{comp} 1,050-1,310 kg/cm²) and a high temperature of deformation under load (the beginning of softening in I, I^P and II takes place at 1,670, 1,540, 1,630°C, respectively, in III at 1,700°C softening did not begin). The content of highly-refractory phases was 86 - 88%. After a storing of 75 days, bricks without burning have also a high deformation temperature (in III there was no deformation at 1,700°C). The test of magnesite-dolomite bricks carried out in the laying of columns of the front wall of 30-t open-hearth furnaces has shown that these bricks are a completely suitable refractory material for them.

V. Zlochevskiy

Card 2/2

131-58-6-8/14

AUTHORS: Kukolev, G. V., Kivin, D. I., Zelenskaya, A. T., Lur'ye, M. A.,
Minskiy, Ya. M.

TITLE: Water-Tight Magnesite-Dolomite Brick (Vodoustoychivyy magnesito-
dolomitovyy kirpich)

PERIODICAL: Ogneupory, 1958, Nr 6, pp. 270 - 274 (USSR)

ABSTRACT: The investigations carried out by the Institute for Refractory
Products showed that by combining magnesite and dolomite in the
raw mixture for clinkers it is possible to obtain products of
high quality, which was proved in the papers by G. V. Kukolev
and D. I. Kivin (Reference 1). In carrying out the present work
clinkers were produced by means of burning a calculated and
controlled finely ground mixture of dolomite, magnesite, quartzite
and phosphorite. The finely ground mixtures were produced accor-
ding to the wet process. In table 1 some results of the labora-
tory investigations are mentioned. In the VNIIO experimental
works several tons of synthetic water-tight magnesit-dolomite
clinkers were produced and of it burned and unburned bricks were
made. Furthermore the production of the masses is described in

Card 1/3

Water- Tight Magnesite-Dolomite Brick

131.58-6-8/14

detail. The investigation of the samples after burning (tables 2 and 3) showed that the bricks of all masses showed a high density and mechanical strength notwithstanding the relatively low burning temperature. In testing the magnesite-dolomite as well as the usual magnesite bricks in practice the former proved to be of better quality. Thanks to the hydraulic hardening the unburned bricks showed after one day of storing a resistance to pressure of 63-83 kg/cm², after one month 294-340 kh/cm², and after 3 months 530-670 kg/cm², having good properties with all this. Furthermore a scheme for the production of magnesite-dolomite bricks is recommended and described in detail. The possibility and usefulness of vacuum filtering of the slip is proved by the work of G. Z. Dolgina (Reference 2). Unburned big magnesite-dolomite blocks can be produced of burned clinker powders in the villages where they are needed. For the metallurgy in the South, Siberia and other districts the production of bricks can be based on the mixture of dolomite and caustic magnesite with additions. These methods are also to be made use for saving magnesite and chromite ores. The production of unburned fire-proof magnesite-dolomite products is to be organized in the works

Card 2/3

Water-Tight Magnesite-Dolomite Brick

131-58-6-8/14

departments for refractory products in the Ural mountains, on the condition that the ready magnesite-dolomite powder of the "Magnesit" will be supplied. Their production of the same burned and unburned products is to be organized in the Nikitovka dolomite Kombinat of dolomite and caustic magnesite with additions. The staff of editors of the periodical remarks on this in reference 3 that first of all a testing of these products of a great industrially produced amount of such bricks would be necessary. There are 3 tables and 2 references, 2 of which are Soviet.

ASSOCIATION: Vsesoyuznyy nauchno-issledovatel'skiy institut ogneporov
(All-Union Scientific Research Institute for Refractories)

1. Refractory materials--Production
2. Refractory materials--Analysis
3. Refractory materials--Test results

Card 3/3

RODDATIS, K.F., kand.tekhn.nauk; KIVINZON, L.M., inzh.

Calculation of the stability of the hydrodynamic characteristics
of the vertical pannels of once-through boilers. Teploenergetika
10 no.1/5-46 Ja '63. (MIRA 16:1)

1. Vsesoyuznyy nauchnyy energeticheskiy institut.
(Boilers)

EYZEN, O.G.; KIVIRYAKHK, S.V.; KOGHEMAN, A.P.; LAUS, T.M.; APPO, I.Kh.

Chemical composition of tar from dictyonemic shale. Khim.i
tekh.topl. 1 masel 5 no.9:37-42 S '60. (MIRA 13:9)

1. Institut khimii AN ESSR.
(Estonia--Oil shale)

KIVISAAR, E.

Treatment of injuries in the abdominal cavity and the inflammation of the
stomach p. 69

SOTSILKTLIK POLIUMJANDUS. POLIUMJANDUS MINISTEERIUM.
Tallin, Hungary. No. 1, 1958.

Monthly List of East European Accessions (EEAI) LC, Vol. 8, no. 11
November 1959.

Uncl.

KIVISAAR, E.

Pine twigs as a vitamin feed for sheep. p. 176.

SOTSIALISTLIK POLLOMAJANDUS. Tallinn, Hungary. Vol. 13, no. 4, Apr. 1958.

Monthly List of East European Accessions (MEAT), *Vol. 8 12 Dec.* LC, No. 4, July, 1959.
Uncl.

~~KIVISML'G. Y. ... (Tallin).~~

large building blocks made of foam slate concrete. Cor. 1 sel'.
stroil. no. 2:6-7 P '57. (MIRA 10:6)
(Tallin--Concrete blocks)

KIVISEL'O, P.P., Cand Tech Sci -- (diss) "Study of ^{under}the
technology and properties of ~~lime~~ shale foamy cement."

Tallin, 1958, 16 pp. (Min of Higher Education USSR.

Tallin Polytechnical Inst) 150 copies (KL, 39-58, 109)

- 37 -

KIVISEL'G, P., kand. tekhn. nauk

Houses built of air-entrained kukermite blocks. Zhil. stroi. no.6:13-15
'59. (MIRA 12:10)

(Estonia--Lightweight concrete) (Estonia--Apartment houses)

KIVISHLG, J.P., insh.

Slate fly-ash concrete in the Estonian S.S.R. Trudy NIIZHB
no.8:106-117 '59. (MIRA 13:4)

1. Institut stroitel'stva i stroitel'nykh materialov AN ESSR.
(Estonia--lightweight concrete)

KIVISELO, Feliks, kand. tekhn. nauk; OJAMAA, Eugen, kand. tekhn.
nauk; IVAND, H., inzh., retsenzent; MASSO, T., red.

[Local building materials] Kohalikud ehitusmaterjalid.
Tallinn, Eesti Riiklik Kirjastus, 1964. 278 p. (In
Estonian)
(MIRA 17:6)

SAKS, E.A.; KIVISEL'O, F.P. [Kiviselg, F.], kand. tekhn. nauk

Panels of shale-ash gas concrete. Stroif. mat. 10 no.6;
29-30 Je '64.

(MCRA 17:10)

1. Zamestitel' nachal'nika Upravleniya promyshlennyykh stroitel'-
nykh materialov Soveta narodnogo khozyaystva Estonskoy SSR (for
Saks).

16.4600

8/023/60/000/002/002/003
0 111/ 0 333

AUTHOR: Kivistik, L.

TITLE: On the Method of Steepest Descent for Solving Non-Linear Equations 16

PERIODICAL: Izvestiya Akademii nauk Estonskoy SSR. Seriya tekhnicheskikh i fiziko-matematicheskikh nauk, 1960, No. 2, pp. 145-159

TEXT: The author considers the equation

$$(1) P(x) = 0,$$

where $P(x)$ is non-linear operator from a real Hilbert space H in H which is twice differentiable in the sense of Frechet. Let the successive approximation of the solution of (1) be carried out according to the scheme

$$(2) x_{n+1} = x_n + \xi_n P(x_n),$$

where ξ_n is the real root of

$$(3) (P(x_n), P(x_n + \xi P(x_n))) = 0.$$

The author proves the convergence of the method under weaker suppositions than Yu. Lumisti (Ref.1) ($P(x)$ need not be a

Card 1/3

83859
8/023/60/000/002/002/003
0 111/ 0 333

On the Method of Steepest Descent for Solving Non-Linear Equations

potential operator) and Guan' Chshaochshi (Ref. 2)

(instead of $(P'(x)h, h) \geq m \cdot \|h\|^2$, $m > 0$, $h \in H$ for all $x \in X$ the same is only demanded for $x = x_0$, where x_0 is the initial approximation).

Then the author considers the modified method according to Altman (Ref.3). He proves convergence in this case too. X

Two theorems deal with the uniqueness of the solution of (1).

8 theorems are given. As an example the author solves an integral equation according to Altman.

L. V. Kantorovich and M. M. Vaynberg are mentioned.

Card 2/3

83859
S/023/60/000/002/002/003
0 111/ 0 333

On the Method of Steepest Descent for Solving Non-Linear Equations

There are 5 references: 4 Soviet and 1 Polish.

ASSOCIATION: Institut energetiki Akademii nauk Estonskoy SSSR
(Institute of Power Engineering of the Academy of
Sciences Estonskaya SSR)

SUBMITTED: June 2, 1959

Card 3/3

8/023/60/000/003/006/012
0111/0222

AUTHOR: Kivistik, L.

TITLE: On Some Iterative Methods for Solving Operator Equations in the Hilbert Space

PERIODICAL: *Investiya Akademii nauk Estonskoy SSR. Seriya Tekhnicheskikh i Fiziko-Matematicheskikh nauk*, 1960, No. 3, pp. 229-241.

TEXT: Let $P(x)$ be an operator two times differentiable according to Frechet, from the real Hilbert space H into the same space. Generalizing the arrangements of Altman (Ref. 1-6) the author proposes the iteration methods

$$(4) \quad x_{n+1} = x_n - \frac{\|P(x_n)\|^2}{\alpha(P'(x_n)y_n, P(x_n))} y_n, \quad n=0, 1, \dots, \quad \frac{1}{2} < \alpha < \infty$$

for the solution of the equation

$$(1) \quad P(x) = 0.$$

Here x_0 is a known initial approximation and either $y_n = \overline{P'(x_n)} P(x_n)$ or $y_n = P(x_n)$, where \overline{P} is the operator conjugated to P . It is proved

Card 1/3

On Some Iterative Methods for Solving
Operator Equations in the Hilbert Space

8/023/60/000/003/006/012
0111/C222

that if $\|P(x_0)\| \leq \delta_0$, $\|P'(x)\| \leq A$, $\|P''(x)\| \leq B$ in a sphere

$\|x - x_0\| \leq r = \frac{M\delta_0}{\alpha(1-q)}$; $\|P'(x)h\| \geq \frac{1}{M}\|h\|$ for all $h \in H$, ($M > 0$) in the same sphere and $q = \frac{1}{\alpha} \sqrt{\alpha^2 - 2\alpha + M^2(A^2 + B\delta_0)} < 1$, then the equation (1) has a solution in the mentioned sphere to which the iteration method (4) $y_n = P'(x_n)P(x_0)$. An estimation for $\|x^* - x_n\|$ is given. Under weaker and stronger assumptions respectively, the author proves a number of further similar assertions. He points to contradictory assumptions in the paper of Altman (Ref.1) (compare Kivistik (Ref.7)). Finally (4) is replaced by the more general arrangement

$$x_{n+1} = x_n - \frac{\|P(x_n)\|^2 y_n}{\alpha_n(P'(x_n)y_n, P(x_n))}, \quad \frac{1}{2} < \alpha_n < \infty.$$

Card 2/3

On Some Iterative Methods for Solving
Operator Equations in the Hilbert Space

S/023/60/000/003/006/012
G111/G222

✓
B

Similar assertions of convergence are given for the new arrangement.
There are 7 references: 1 Soviet and 6 Polish.

ASSOCIATION: Institut energetiki Akademii nauk Estonskoy SSR (Power
Engineering Institute of the Academy of Sciences of the
Estonian SSR)

SUBMITTED: November 2, 1959

Card 3/3

21311

S/023/60/000/004/001/005
D221/D305

16.6500

AUTHOR: Kivistik, L.

TITLE: One generalization of Newton's method of approximation

PERIODICAL: Akademiya nauk Estonskoy SSR. Izvestiya. Seriya fiziko-matematicheskikh i tekhnicheskikh nauk, no. 4, 1960, 301-312

TEXT: Let $P(x) = 0$ (1)

be a non-linear operator in the Banach space X into the space Y of the same type. To solve this equation the author considers two iterated methods, based on

$$x_{n+1} = x_n - \alpha_n \Gamma(x_n) P(x_n) \quad (2)$$

where $\Gamma(x) = [P'(x)]^{-1}$ and α_n are numbers within the interval (0.2) with some additional limitations. When $\alpha_n = 1$ ($n = 0, 1, \dots$)

Card 1/16

21311

S /023/60/000/004/001/005
D221/D305

One generalization of ...

then Eq. (2) gives the well known Newton's method of approximation; if $0 < \alpha_n = \alpha < 1$ then Eq. (2) resolves into the generalized method of D.A. Grave as cited by L.V. Kantorovich (Ref. 1: Metode N'yutona (On the Newton Method), Tr. Matem. in-Ta im V.A. Steklova, 28, 1949, 104-144) /Abstractor's note: Surname Grave transliterated from Russian/. The congruence of the two methods is proved by the author with the help of theorems established by Kantorovich (Ref. 1: Op.cit.). First, a subsidiary theorem A is established. Let the following conditions be satisfied: 1) There exists an inverse operator $\Gamma(X_0) = [P'(X_0)]^{-1}$ and also $\|\Gamma(X_0)\| \leq B_0$; 2) $\|\Gamma(X_0)P(X_0)\| \leq \eta_0$; 3) for all $X \in S(X_0, r)$, where the symbol $S(X_0, r)$ denotes the sphere $\|X - X_0\| \leq r$ and r is given by

$$r = N(h_0) \eta_0 = \frac{1 - \sqrt{1 - 2h_0}}{h_0} \eta_0$$

Card 2/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

the following inequality is satisfied $\|P^*(X)\| \leq K_1 \cdot 4) h_0 = B_0 K \eta_0 \leq \frac{1}{2}$. Then equation $P(X) = 0$ has within the sphere $S(X_0, r)$ the solution X^* , to which converge successive approximations of Eq. (2), where $0 < \alpha_n = \alpha < 1$. The theorems proper, proving the convergence of Eq. (1) follow next. Theorem 1. Let conditions 1-4 of theorem A be satisfied and $0 < \alpha_n < 1$, $\alpha = \inf \alpha_n > 0$, then equation $P(X) = 0$ has within the sphere $S(X_0, r)$, the solution X^* , to which converge consecutive $\{X_n\}$ obtained from (1) and the following are the error estimates

$$\|X^* - X_n\| \leq N(\bar{h}_n) \|\Gamma(X_n) P(X_n)\| \leq N(h_0) \eta_0 \cdot q^n, \quad (6)$$

where

$$\bar{h}_n = \|\Gamma(X_n)\| K \|\Gamma(X_n) P(X_n)\| \text{ and } q = \frac{1 - \alpha + \frac{1}{2} \alpha^2 h_0}{1 - \alpha h_0}.$$

Card 3/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

Theorem 2. If conditions 1-4 of theorem A are satisfied and numbers α_n chosen to satisfy

$$1 > \alpha_n > \frac{1 - \sqrt{1 - 2h_n(1 - \gamma h_n)}}{h_n}, \quad (21)$$

and also

$$\frac{1}{2} < \gamma < \frac{1 - h_0}{N(h_0)h_0} \quad (22)$$

is satisfied, then the error in approximations obtained by using Eq. (1) can be evaluated from formula

$$\|x^* - x_n\| \leq N(h_0)\eta_0(1 - h_0)^n \left[\frac{N(h_0)h_0}{1 - h_0} \right]^{2^{n-1}} \quad (20)$$

or even more accurately from

Card 4/16

2311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

$$\begin{aligned} \|x^* - x_n\| &\leq N(h_n) \eta_n \leq N(h_n) \frac{\eta_0 \cdot \gamma^{h_{n-1}} \dots \gamma^{h_1} \cdot \gamma^{h_0}}{(1 - \alpha_{n-1} h_{n-1}) \dots (1 - \alpha_0 h_0)} < \\ &\leq \frac{N(h_0) \eta_0 \lfloor \gamma^{N(h_0) h_0} \rfloor^{2^{n-1}}}{(1 - \alpha_{n-2} h_{n-2})^{2^1 - 1} \dots (1 - \alpha_0 h_0)^{2^{n-1} - 1}} \leq \end{aligned} \quad (19)$$

$$\leq \frac{N^2(h_0)}{N(h_n)} \eta_0 (1 - \alpha_{n-1} h_{n-1}) \dots (1 - \alpha_0 h_0) \lfloor \gamma^{N^2(h_0) h_0} \rfloor^{2^{n-1}}.$$

In the latter it is enough, instead of inequality (22), to satisfy the condition $\frac{1}{2} \leq \gamma \leq \lfloor N^2(h_0) h_0 \rfloor^{-1}$.

Card 5/16

21311

S/023/60/000/004/001/005

D221/D305

One generalization of ...

Theorem 3. follows from Theorem 2. If conditions 1-4 of Theorem A are satisfied, the error in approximations, obtained by Newton's method can be evaluated from

$$\begin{aligned} \|x^* - x_n\| &\leq N(\bar{h}_n) \|f(x_n) P(x_n)\| \leq N(h_n) \eta_n = \\ &= \frac{N^2(h_n)}{N(h_n)} \eta_n \left[\prod_{k=0}^{n-1} (1 - h_k) \right] \cdot \left[\frac{1}{2} N^2(h_0) h_0 \right]^{n-1} < \\ &< N(h_0) \eta_n (1 - h_0)^n \left[\frac{N(h_0) h_0}{2(1 - h_0)} \right]^{n-1}, \end{aligned}$$

where $\bar{h}_n = \|f(x_n)\|/K/\|f(x_n) P(x_n)\|$ and h_k ($k = 1, 2, \dots$) are determined recurrently by formula

$$\begin{aligned} \eta_{n+1} &= \frac{1}{2} \frac{h_n}{1 - h_n} \eta_n, \\ h_{n+1} &= \frac{1}{2} \frac{h_n^2}{(1 - h_n)^2}. \end{aligned} \quad (23)$$

Card 6/16

21311

S/023/00/000/004/001/005
D221/D305

One generalization of ...

Theorem 3 for $h_0 < \frac{1}{2}$ gives a better error estimate than that obtained using the Kantorovich (Ref. 1; Op.cit.) method or by S.Yu. U'lm (Ref. 2: O skhodimosti nekotorykh iteratsionnykh protsessov v prostranstve Banakha (On the Convergence of some Iterative Processes in the Banach Space), Uch. zap. Tartusk. Gos. un-ta, 42, 1956, 135-142). Theorem 4. Let the following conditions be satisfied. 1) There exists an inverse operator $\Gamma(X_0) = [P'(X_0)]^{-1}$ with $\|\Gamma(X_0)\| \leq B_0$; 2) $\|\Gamma(X_0) P(X_0)\| \leq \eta_0$; 3) for all $X \in S(X_0, r)$, where

$$r = \frac{B\eta_0}{1-q} \text{ and } q = \max \left\{ \frac{1 - \alpha + \frac{1}{2}\alpha^2 h_0}{1 - \alpha h_0}, \right.$$

$$\left. \frac{B - 1 + \frac{1}{2} B^2 h_0}{1 - B h_0} \right\}$$

Card 7/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

there is an estimate $\|P^n(X)\| \leq K$; 4) The inequalities

$$\|1 - \alpha\| + \frac{1}{2} \alpha^2 h_0 \leq (1 - \alpha h_0)^2,$$

and

$$\beta - 1 + \frac{1}{2} \beta^2 h_0 \leq (1 - \beta h_0)^2, \beta h_0 < 1,$$

where $h_0 = B_0 K_{\gamma_0}$; Equation $P(X) = 0$ then has within the sphere $S(X_0, r)$ the solution x^* , to which converge the consecutive approximations X_n as obtained from Eq.(2), where $0 < \alpha_n < 2$ and the error estimate becomes

$$\|x^* - X_n\| \leq \frac{\beta}{1-q} \|\Gamma(X_n)\| \|P(X_n)\| \leq \frac{\beta_{\gamma_0}}{1-q} q^n. \quad (24)$$

Let the limit of the norm of operator $\Gamma(x)$ now be known in the

Card 8/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

whole of the domain (Ref. 3: I.P. Mysovskikh, K voprosu O skhodimosti metoda N'yutona (On the Convergence of the Newton Method), Tr. Matem., in-ta im. V.A. Steklova, 28, 1949, 145-147), and (Ref. 4: O skhodimosti metoda L.V. Kantorovicha dlya resheniya nelineynykh funktsional'nykh uravneniy i ego primeneniya (On the Convergence of the L.V. Kantorovich method for Resolving Non-Linear Functional Equations and its Uses), Vestn. Leningr. un-ta, no. 11, 1953, 25-48). If $0 < \alpha_n \leq 1$, $\alpha = \inf \alpha_n > 0$, $\beta = \sup \alpha_n$ then the following Theorem 5 holds: Let the following conditions be satisfied: 1) $\|P(X_0)\| \leq \delta_0$; 2) For all $X \in S(X_0, r)$, where

$$r = \frac{\beta \delta_0}{1 - \alpha},$$

there exists the operator $\Gamma(X)$ with $\|\Gamma(X)\| \leq B$. 3) For all $X \in S(X_0, r)$ $\|P''(X)\| \leq K$; 4) $\beta h_0 = \beta B^2 K \delta_0 < 2$. Then $P(X) = 0$ has

Card 9/16

21311
S/023/60/000/004/001/005
D221/D305

One generalization of ...

within the sphere $S(X_0, r)$ the solution X^* , to which converge the consecutive approximations X_n obtained from Eq. (2) and the following estimate exists.

$$\|X^* - X_n\| \leq \frac{B\delta}{1-q} \|P(X_n)\| \leq \frac{B\delta_0}{1-q} \cdot q^n,$$

where $q = \max \left\{ 1 - \alpha + \frac{1}{2} \alpha^2 h_0, 1 - B + \frac{1}{2} B^2 h_0 \right\}$.

Let conditions 2 and 3 of Theorem 5 be satisfied for all elements X of a certain sphere $S(X_0, R)$. Making α small enough it may always be that $\alpha h_0 < 2$ and $r = r(\alpha) < R$ if only $R > B\delta_0$. The next theorem can, therefore, be formulated as Theorem 6. If within a certain sphere $S(X_0, R)$ where $R > B \|P(X_0)\|$, there exists the inverse operator $T(X) = [P'(X)]^{-1}$ the norm of which is limited by

Card 10/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

number $B // \Gamma(X) // < B$ and $//P^*(X)//$ is also limited within sphere $S(X_0, R)$, then $P(X) = 0$ has a solution within this sphere and also the converging to this solution. Approximations can be determined from Eq. (2) where $\alpha_n = \alpha$ and $\alpha =$ a sufficiently small positive number. A more generalized method of evaluation follows given as

$$X_{n+1} = X_n - A_n \Gamma(X_n) P(X_n), \quad (25)$$

where A_n are arbitrary linear operators from space X into the same space. In particular if $A_n = \alpha_n E$, where $\alpha_n \in (0, 2)$ - the method of Eq. (2) is obtained. Using the Taylor formula and identity

$$P(X_n) = P'(X_n) \Gamma(X_n) P(X_n)$$

and

$$P(X_{n+1}) = P'(X_n)(E - A_n) \Gamma(X_n) P(X_n) + R_n \quad (26)$$

Card 11/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

is obtained from which

$$\Gamma(X_{n+1}) P(X_{n+1}) = H_n (E - A_n) I(X_n) P(X_n) + \Gamma(X_{n+1}) R_n \quad (29)$$

follows, where

$$\|R_n\| \leq \frac{1}{2} \|P''(X_n + \tau_n(X_{n+1} - X_n))\| \|X_{n+1} - X_n\|^2, \quad 0 \leq \tau_n \leq 1$$

and $H_n = [E - \Gamma(X_n)(P'(X_n) - P'(X_{n+1}))]^{-1}$. Using the identity (29) and the methods of deducing previous theorems the following theorems can be obtained: Theorem 7. Let the following conditions be satisfied: 1) There exists the inverse operator $\Gamma(X_0) = [P'(X_0)]^{-1}$, with $\|\Gamma(X_0)\| \leq B_0$; 2) $\|\Gamma(X_0) P(X_0)\| \leq \eta_0$;

3) For all $x \in S(X_0, r)$, where $r = \frac{b\eta_0}{1-q}$, there is an error esti-

Card 12/16

21311
S/023/60/000/004/001/005
D221/D305

One generalization of ...

mate $\|P^n(X)\| \leq K$; 4) The inequality $e + \frac{1}{2} b^2 h_0 < (1 - b h_0)^2$ is

satisfied, where $b h_0 = b B_0 K \eta_0 < 1$, $b = \sup \alpha_n < 2$, $q = \frac{e + \frac{1}{2} b^2 h_0}{1 - b h_0}$,

and α_n and e are determined from

$$\|E - A_n\| \leq e_n, \quad \|A_n\| \leq \alpha_n, \quad (26)$$

and

$$e_n \leq e < 1. \quad (27)$$

Then $P(X) = 0$ has within the sphere $S(X_0, r)$ the solution X^* , to which converge the consecutive approximations X_n obtained from Eq. (25) and the error estimates become as given by

$$\|X^* - X_n\| \leq \frac{b}{1-q} \| \Gamma(X_n) P(X_n) \| \leq \frac{b \eta_0}{1-q} \cdot q^n.$$

Card 13/16.

21311
S/023/60/000/004/001/005
D221/D305

One generalization of ...

Theorem 8. Let the following conditions be satisfied: 1) There exists the inverse operator $\Gamma(X_0) = [P'(X_0)]^{-1}$, with $\|\Gamma(X_0)\|/K \leq B_0$; 2) $\|\Gamma(X_0) P(X_0)\| \leq \eta_0$; 3) For all $x \in S(X_0, r)$, where

$r = \frac{b\eta_0}{1 - (1 - bh_0)I_0}$ the inequality $\|P^n(X)\| \leq K$ is satisfied;

4) The operators A_n are chosen so that for all n $e_n + \frac{1}{2}a_n^2 h_n \leq \gamma h_n$ holds, where γ satisfies the condition

$\frac{1}{2} \leq \gamma \leq \frac{(1 - bh_0)^2}{h_0}$ and $bh_0 = bB_0 K \eta_0 < 1$ and $h_n (n > 1)$ are deter-

mined by the recurrent relationship

$$h_n = \frac{e_{n-1} + \frac{1}{2}a_{n-1}^2 h_{n-1}}{(1 - e_{n-1} h_{n-1})^2} h_{n-1}.$$

Card 14/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

Then $P(X) = 0$ has within the sphere $S(X_0, r)$ the solution X^* , to which converge the consecutive approximations X_n obtained from (25) and the error evaluations become

$$\|X^* - X_n\| \leq \frac{b\eta_0 (1 - b\eta_0)^n}{1 - (1 - b\eta_0) l_0^{2^n}} \cdot l_0^{2^n - 1},$$

where $l_0 = \frac{\eta_0}{(1 - b\eta_0)^2}$ ($b = \sup \alpha_n$). The application of (2) and of (25) may be considered as an approximate evaluation of the consecutive approximation of $P(X) = 0$ using the Newton Method.

$$X_{n+1} = X_n - \Gamma(X_n) P(X_n) \quad (30)$$

thus, if consecutive approximations X_{n+1} are determined in the Card 15/16

21311

S/023/60/000/004/001/005
D221/D305

One generalization of ...

sense that instead of the increment - $\Gamma(X_n) P(X_n)$ a certain other element - $\alpha_n \Gamma(X_n) P(X_n)$ or $\Lambda_n \Gamma(X_n) P(X_n)$ is found, in which $\alpha_n \in (0, 2)$ or $\|E - \Lambda_n\| \leq \epsilon < 1$ respectively, then the obtained approximations still converge towards the exact solutions (Theorems 1, 4, 5, 6, 7). If the permissible error is not too great the convergence will still remain that of the second order (Theorems 2, 8).
/Abstractor's note: No definitions of symbols are given by the author. Although it is not mentioned explicitly - all symbolic notation seems to follow that used by L.V. Kantorovich in Ref. 1: Op. cit.]. There are 6 Soviet-bloc references.

ASSOCIATION: Institut kibernetiki akademii nauk Estonskoy SSR
(Institute of Cybernetics of the Academy of Sciences of Estonian SSR).

SUBMITTED: February 10, 1960

Card 16/16

88557

16.4600

16.6500

8/020/61/136/001/002/037
C111/C222

AUTHOR: Kivistik, L.A.

TITLE: On a Modification of the Iterative Method With Minimal Residuals
for the Solution of Nonlinear Operator Equations

PERIODICAL: Doklady Akademii nauk SSSR, 1961, Vol. 136, No. 1, pp.22-25

TEXT: Let $P(x)$ be an operator from H into H which is two times differentiable according to Frechet, let H be a real Hilbert space. The solution of

$$(1) \quad P(x) = 0$$

is carried out with the arrangement

$$(2) \quad x_{n+1} = x_n + \epsilon_n y_n, \quad n = 0, 1, \dots$$

where $x_0 \in H$ is the initial approximation, ϵ_n is chosen so that for a fixed y_n the expression $\|P(x_n) + P'(x_n)(x_{n+1} - x_n)\|^2 = \|P(x_n) + \epsilon_n P'(x_n)y_n\|^2$ becomes minimal.

Card 1/6

88557

9/020/61/136/001/002/037
C111/C222

On a Modification of the Iterative Method With Minimal Residuals for the Solution of Nonlinear Operator Equations

Let $y_n = P(x_n)$. Then one obtains the method

$$(6) \quad x_{n+1} = x_n - \frac{(P(x_n), P'(x_n)P(x_n))}{\|P'(x_n)P(x_n)\|^2} P(x_n)$$

Theorem 1: Let the following conditions be satisfied :

1° $\|P(x_0)\| \leq \delta_0$

2° For all $x \in S(x_0, r)$, where $r = \frac{M\delta_0}{1-q}$, and $S(x_0, r)$ denotes the sphere $\|x - x_0\| \leq r$ let :

a) $\|P'(x)\| \leq A$ b) $\|P''(x)\| \leq B$ c) $|(P'(x)h, h)| \geq M^{-1}\|h\|^2$ for all $h \in H (M > 0)$

3° $q = \sqrt{1-b^{-1}} + \frac{1}{2} a_0 < 1$, where $b = M^2 A^2$, $a_0 = M^2 B \delta_0$.

Then (1) has a unique solution x^* in $S(x_0, r)$ to which there converges
Card 2/6

08557

S/020/61/136/001/002/037
C111/C222

On a Modification of the Iterative Method With Minimal Residuals for the Solution of Nonlinear Operator Equations

the sequence $\{x_n\}$ obtained by (6), where

$$(7) \quad \|x^* - x_n\| \leq M \|P(x_n)\| \leq M \delta_0 q^n$$

Theorem 2 : Let

$$1^\circ \quad \|P(x_0)\| = \delta_0 \leq \bar{\delta}_0$$

$$2^\circ \quad |(P'(x_0)h, h)| \geq M_0^{-1} \|h\|^2 \text{ for all } h \in H (M_0 > 0).$$

$$3^\circ \quad \text{Let } \|P'(x)\| \leq A, \quad \|P''(x)\| \leq B \text{ be valid for all } x \in S(x_0, r),$$

where $r = \frac{1}{B} \left(\frac{1}{M_0} - \frac{1}{M} \right) \frac{\delta_0}{\bar{\delta}_0} (M^+ = \lim M_n \leq +\infty)$.

$$4^\circ \quad \text{Let the magnitudes } a_0 = M_0^2 B \bar{\delta}_0 \text{ and } b_0 = M_0^2 A^2 \text{ be so that the}$$

sequence $\{a_n\} = \{M_n^2 B \bar{\delta}_n\}$ calculated with the aid of the recurrence formulas

Card 3/6

88557

S/020/61/136/001/002/037
C111/G222

On a Modification of the Iterative Method With Minimal Residuals for the Solution of Nonlinear Operator Equations

$$(8) \quad \begin{aligned} M_{k+1} &= \frac{M_k}{1 - M_k^2 B \bar{\delta}_k} \\ \bar{\delta}_{k+1} &= \bar{\delta}_k \left(\sqrt{1 - (M_k^2 A^2)^{-1}} + \frac{1}{2} M_k^2 B \bar{\delta}_k \right) \end{aligned}$$

is convergent (i.e. that $a_n < 1$ for all n).

Then (1) has a solution x^* in $S(x_0, r)$ to which there converges the sequence $\{x_n\}$ obtained by (6), and here it is

$$(9) \quad \|x^* - x_n\| \leq \frac{2M_n \bar{\delta}_n}{1 + \sqrt{1 - 2M_n^2 B \bar{\delta}_n}} < 2M_n \bar{\delta}_n$$

where $\bar{\delta}_n = \|P(x_n)\|$ and M_n is determined according to (8). If $M^f < \infty$ or $\bar{\delta}_0 > \bar{\delta}_0$ then the solution is unique in $S(x_0, r)$.

Card 4/6

88557

S/020/61/136/001/002/037

G111/C222

On a Modification of the Iterative Method With Minimal Residuals for the Solution of Nonlinear Operator Equations

Theorem 3 : If $a_{00} b_{00} \leq \frac{1}{g}$ then the condition 4⁰ of theorem 2 is satisfied.

Choosing $y_n = \overline{P'(x_n)} P(x_n)$, where $\overline{P'(x)}$ is adjoint to the linear operator $P'(x)$ then one obtains the method

$$(10) \quad x_{n+1} = x_n - \frac{\| \overline{P'(x_n)} P(x_n) \|^2}{\| P'(x_n) \overline{P'(x_n)} P(x_n) \|^2} \overline{P'(x_n)} P(x_n)$$

Theorem 5 : Let the conditions of theorem 2 be satisfied with the exception of the condition 2⁰ and the relations (8) which are replaced by the condition

$\| P'(x_0)h \| \geq M_0^{-1} \| h \|$ and $\| \overline{P'(x_0)}h \| \geq M_0^{-1} \| h \|$ for all $h \in H$ ($M_0 > 0$) and

the relations

Card 5/6

00557

S/020/61/136/001/002/037
C111/C222

On a Modification of the Iterative Method With Minimal Residuals for the Solution of Nonlinear Operator Equations

$$(11) \quad M_{k+1} = \frac{M_k}{1 - M_k^2 B \bar{S}_k}, \quad \bar{S}_{k+1} = \bar{S}_k \left(\frac{M_k^2 A^2 - 1}{M_k^2 A^2 + 1} + \frac{1}{2} M_k^2 B \bar{S}_k \right)$$

Then (1) has a solution x^* in $S(x_0, r)$ to which there converges the sequence $\{x_n\}$ obtained with the aid of (10), and there hold the estimations (9), where $\bar{S}_n = \|P(x_n)\|$, and M_n are calculated according to (11).

Theorem 6 : If $(b_0 + 1)(g - 12a_0 + 8a_0^2 - 2a_0^3)a_0 \leq 4$ and $a_0 \leq 4/g$, then the condition 4° of theorem 5 is satisfied.

The author mentions M.A. Krasnosel'skiy and S.G. Kreyn. There are 3 references : 2 Soviet and 1 American.

ASSOCIATION: Institut energetiki Akademii nauk Estonskoy SSR (Power Engineering Institute of the Academy of Sciences Estonskaya SSR)

PRESENTED: July 15, 1960, by S.L. Sobolev, Academician

SUBMITTED: June 14, 1960

Card 6/6

KIVISTIK, L. A.

Cand Phys-Math Sci - (diss) "Iteration methods in Hilbert space."
Tartu, 1961. 10 pp; (Tartu State Univ); 250 copies; free;
bibliography at end of text (15 entries); (KL, 5-61 sup, 173)

KIVISTIK, L.A.

~~Approximation~~ iterative method with minimal residues for the
solution of nonlinear operator equations. Dokl.AN SSSR 136 no.1:
22-25 Ja '61. (MIRA 14:5)

1. Institut energetiki Akademii nauk Estonskoy SSR. Predstavleno
akademikom S.L.Sobolevym.
(Differential equations, Partial) (Operators (Mathematics))

A centrifugal plant for the thermal conversion of oil shale ores. N. H. Kuznetsov, N. I. Kishchik, B. I. Tyagunov, and A. A. Klevt. *Chemical Abstracts*, 1956, No. 5, 6-18. — An experimental plant for working up 830 tons/day of ore grade oil shale ores, containing approx. 6% moisture, 45% ash, 15% C, and 80% org. matter, operates at 750 to 800°. Liquid products representing 10% of the dry wt. of the shale consist of 15-21% light oils, 45-75% heavy oils, and 8-27% tars. Gas yields with thermal values from 10,000 to 12,000 kcal/cu. m. are from 35 to 45 cu. m./ton. The gas composition is CO_2 8.5, CO 12.0, C_2H_4 0.2, C_2H_6 7.6, C_3H_8 8.9, C_4H_{10} 4.3, C_4H_8 1.1, CO_2 0.3, H, 7.8, CH, 16.7, N, 10.8, and C_2 and higher 3.1%. The ash shows promise as a raw material for making cements of the so-called "redmen" type. On the basis of the exptl. results, plans are being drawn up for the construction of a full-scale plant with 1,000 tons/day capacity.

KIVII, A.A.

AARNA, A.Ya. [Aarna, A.J.], doktor tekhnicheskikh nauk, retsentsent; KULL', M. [Kull, M.], kandidat ekonomicheskikh nauk, retsentsent; KYLL', A.F. [Kyll, A.F.], redaktor; KIVIT, A.A., redaktor; MIKHELIS, K.A. [Mihelis, K.A.], redaktor; GUBERGILIS, Mark Yakovlevich, redaktor; ROGINA, G.M., vedushchiy redaktor; YASHCHURZHINSKAYA, A.B., tekhnicheskiiy redaktor

[Engineering and economic problems of industrial semicoking of combustible shale; a collection of papers] Voprosy tekhniki i ekonomiki promyshlennogo polukoksovaniia goriuchikh slantssev; sbornik statei. Leningrad, Gos.nauchno-tekhn. izd-vo neft. i gorno-toplivnoi lit-ry, Leningr.otsd-nie, 1957. 337 p. (MLRA 10:7)

1. Kivioli Polevkiiviseia Kombinsat.
(Oil shales)

XIVIT, A.A., red.; ANTONS, R.I., red.; AARNA, A.Ya., prof., doktor
tekhn.nauk, retsentsent; KUL'L', B.V., kand.ekon.nauk, retsentsent;
RAZINA, G.M., vedushchiy red.; YASHCHURZHINSKAYA, A.B., tekhn.red.

[Technology and economic aspects of the industrial semicoking of
oil shales] Voprosy tekhniki i ekonomiki promyshlennogo polukokso-
vaniia goruchikh slantsev. Leningrad, Gos.nauchno-tekhn.isd-vo
neft. i gorno-toplivnoi lit-ry, Leningr.otd-nie. No.2. 1959.
429 p. (MIRA 12:10)

1. Kiviõli Põlevkiviseemia Kombinaat.
(Oil shales)

KUZNETSOV, Dmitriy Trofimovich; ZHUKOV, V.A., dotsent, retsenzent;
KIVIT, A.A., nauchnyy red.; NIKOLAYEV, G.S., nauchnyy red.;
KUDINA, G.W., vedushchiy red.; YASHCHURZHINSKAYA, A.B.,
tekhn.red.

[Outline of the development of the oil-shale industry in the
 Estonian S.S.R.] Ocherki razvitiia slantsevoi promyshlennosti
 Estonskoi SSR. Leningrad, Gos.nauchno-tekhn.isd-vo neft. i
 gorno-toplivnoi lit-ry, Leningradskoe otd-nie, 1960. 199 p.
 (MIRA 13:6)

1. Zaveduyushchiy kafedroy khimicheskoy tekhnologii Leningradskogo
 inzhenerno-ekonomicheskogo instituta (for Zhukov).
 (Estonia--Oil shales)

KIVIVYALI, B.T.

Manufacturing skis of plywood. Der.prom. 8 no.6:24-25
Je '59. (MIRA 12:8)

1. Tallinskaya lyzhnaya fabrika "Dinamo."
(Skis and skiing)

PEKHOM, K.M. [Pehme, K.]; KIVIVIALI, B.T. [Kiviväli, B.]

Design and technology for the manufacture of skis from
plywood. Der.prom. 9 no.3:21 Nr '60.
(MIRA 13:6)

1. Lyskymäsa fabrika Neitonskogo respublikanskogo soveta
"Dinamo".

(Neitonia—Skis and skiing)

KIVIVYALI, B.T. [Kiviväli, B.T.]; KEROE, R.H. [Kärde, R.H.]

Efficient cutting of logs for the manufacture of multiple layer
skis. Der.prom. 9 no.9:19-21 8 '60. (MIRA 13:9)
(Matonia--Skis and skiing) (Woodwork)

KIVKIND, A. I. (Kazan)

"Proton Resonance in Paramagnetic solutions, paper presented at the International Conference on Physics of Magnetic Phenomena, Overdlovsk, USSR, 23-31 May 1956.

KIVKUTJAN, F. R.

USSR/Chemistry - Biochemistry

Card 1/1 : Pub. 22 - 29/44

Authors : Sisakyan, N. H., Memb. Corresp. of Acad. of So. USSR; Besinger, E. H.,
and Kivkutsan, F. R.

Title : Amino-acid composition of phycoerythrin

Periodical : Dok. AN SSSR 98/1, 111-114, Sep 1, 1954

Abstract : The amino-acid composition of phycoerythrin (chromoprotein) derived
from *Callithamnion rubescens* algae and containing 10.61% of N per
dry weight, was investigated. The amino-acids were identified by
the method of distributive chromatography on paper. The results
obtained are shown in tables. Sixteen references: 8-USA; 3-German
and 5-USSR (1928-1954). Tables; drawings; illustrations.

Institution : Acad. of So. USSR, The A. N. Bakh Institute of Biochemistry

Submitted : June 11, 1954

KITMAN, A.M.

Pneumatic high-frequency drilling hammers. Biol.tekh.-ekon.inform.
no.7:6-7 '58. (MIRA 11:9)
(Boring machinery)

KIVMAN, A.M.

The 10-10 pneumatic breaking hammer. Biul.tekh.-ekon.inform. no. 48
7-8 '99. (MIRA 12:7)
(Coal mining machinery)

KIVMAN, A.M., inzh.

Select the best models of air drills. Shakht. stroi. 7 no.2:6-9
F '63. (MIRA 16'3)

1. Zavod "Pnevmatika", Leningrad.
(Boring machinery—Pneumatic driving)

KIVMAN, P.M.

Control device for the measurement of forging shifts. Kus.-
shtam.proizv. 4 no.8147 Ag 162. (MIRA 1518)
(Forging machines--Attachments)

KIVNAN, G. Ya., SHITKOV, I. L. and GILBOVITSKIY, Ye. B.

"Apparatus for the Graphic Recording of Changes in the Lumen of Blood Vessels During Physiological Experiment." *Farmakol i Toksikol*, No. 2, p 57, 1951.

KIVMAN, G. YA.

USSR/Medicine, Biology - Antibiotics - Jul/Aug 72

"Control of Streptomycin in Regard to Toxicity and Pyrogenicity," A. Ye. Tsyakina, G. Ya. Kivman, Control Inst of Sera and Vaccines imeni L. A. Tarasovich

"Med Prom" No 4, pp 42-44

Describes in detail procedures for testing toxicity on mice and pyrogenicity on rabbits in order to control the quality of streptomycin.

22279

1. KIVMAN, G. Ya
2. USSR (600)
4. Antibiotics
7. Antibiotics as stimulants of growth in animals, Antibiotiki, 5, No. 4, 1952.

9. Monthly List of Russian Accessions, Library of Congress, February 1953, Unclassified.

TEBYAKINA, A.Ye.; SHIRYAYOVA, V.L.; KIVMAN, G.Ye.

Comparative study of methods of determining the activity of penicillin.
Trudy VNIIA no.1:97-104 '53. (MLA 8:1)

1. In Tsentral'nogo gosudarstvennogo nauchnogo kontrol'nogo instituta
im. Tarasevicha (direktor S.I.Didenko).
(Penicillin)

KIVMAN, G. Ya.

Antibiotics

Allergy to antibiotics. Antibiotiki. 6, No. 1, 1953.

9. Monthly List of Russian Accessions, Library of Congress, June 1953, Uncl.

KIVMAN, O.Ya., kandidat meditsinskikh nauk.

Effect of antibiotics on the growth of animals. Priroda 42 no.11:98-100 N
'53. (MIRA 6:11)

1. Gosudarstvennyy kontrol'nyy institut im. L.A.Tarasovicha.
(Antibiotics) (Growth)

KIVMAN, O.Ya., kandidat meditsinskikh nauk.

Effect of antibiotics upon the protective reactions in the body.

Antibiotiki 7 no.2:14-27 '54.

(MLRA 7:4)

(Antibiotics)

KIVMAN, O.Ya. (Moskva)

Effect of antibiotics on growth of animals and plants. Usp. sov. biol. 38 no.2:163-182 8-0 '54. (MLA 8:1)

(ANTIBIOTICS, effects,
on growth of animals & plants)
(PLANTS, effect of drugs on,
antibiotics, on growth)
(GROWTH, effect of drugs on,
antibiotics, in animals & plants)

KIVMAN, G. YA.
USSR/Biochemistry

Card 1/1

Authors : Sisakyan, N. M; Besinger, E. N; Oarkavi, P. G., and Kivman, G. Ya.

Title : Simple method determining amino-acids through chromatographic analysis on paper.

Periodical : Dokl. AN SSSR, 96, Bd. 2, 343 - 346, May 1954

Abstract : Determination of amino-acids is carried out by a two-dimensional chromatographic method. The initial process of separation is accomplished with the aid of methyl alcohol - water - pyridine (10 : 10 : 2) and the second and final process with n-butyl alcohol - methylethylketone - water - diethylamine (20 : 20 : 10 : 2). The solvents require no preliminary purification. The appearance of amino-acid on the paper is attained by treating the latter in a 0.4-% ninhydrin solution in methyl alcohol, in acetone or n-butyl alcohol. One reference. Table, photos.

Institution : Acad. of Sci. USSR, The A. N. Bekh Inst. of Biochemistry and the State Control Inst. of Serums and Vaccines at the Ministry of Health USSR.

Submitted : March 13, 1954

MIN translation in /M Inclusion - B-93873, 28 Nov 55

TRBYANKINA, A.Ye.; KIVMAN, O.Ya.; SVIRSKAYA, S.I.

Pharmacological control of streptomycin. Med.prom. no.1:23-26 Ja-Mr
'55. (MIRA 8:5)

1. Kontrol'nyy institut syverotok i vaksain imeni Tarasevicha.
(STREPTOMYCIN,
standard)

KIVMAN, G. Ya.

KIVMAN, G.Ya., kandidat meditsinskikh nauk

Effect of antibiotics on the liver in men and animals. Antibiotiki
8 no.1:19-24 '55. (MLA 8:3)

(LIVER, effect of drugs on,
antibiotics)

(ANTIBIOTICS, effects,
on liver)

KIVMAN, O.Ya., kandidat meditsinskikh nauk; KHASKIN, L.S.

Utilisation and sterilisation of side products obtained during
production of antibiotics; review of foreign periodical literature.

Antibiotiki 8 no.2:25-36 '55.

(MIRA 8:5)

(ANTIBIOTICS, preparation of,
use of side products, review)

(DRUG INDUSTRY,
use of side products in antibiotic indust., review)

KIVMAN, G.Ya.; PRYADKINA, M.D.; GUTCHOVA, N.M.

Preparation for the detection and stimulation of growth of *Vibrio* comma. Zhur. mikrobiol. epid. i immun. no.12:61-66 D '55. (MIRA 9:5)

1. Is laboratorii Gosudarstvennogo kontrol'nogo instituta syvorotok i vaktsin imeni L.A. Tarashevicha (dir.-S.I. Didenko)

(*VIBRIO COMMA*, culture,

medium containing *Bacillus mesentericus* filtrates
for detection & stimulation of growth.)

(*BACILLUS*,

mesentericus, filtrates in culture media for detection
& stimulation of growth of *Vibrio comma*)

(*CULTURE MEDIA*,

for *Vibrio comma*, eff. of *Bacillus mesentericus* filtrates
in detection & stimulation of growth)

KIVMAN, O.Ya.

Dynamics of the binding of tetracyclines by the liver. Antibiotiki
1 no.3:42-46 My-Je '56. (MIRA 9:10)

1. Otdel eksperimental'noy khimioterapii (sav. ohlen-korrespondent
AMN SSSR prof. Kh.Kh.Planel'yes) Instituta farmakologii i eksperi-
mental'noy khimioterapii AMN SSSR

(LIVER, metabolism,
tetracycline, binding (Rus))

(TETRACYCLINE, metabolism,
liver, binding (Rus))

"Second Volume of Works on Biomyxin," by G. Ya. Kivman,
Antibiotiki, Vol 1, No 9, Sep/Oct 56, pp 59-60

This work reviews a collection of articles on the antibiotic, biomyxin issued under the editorship of Z. V. Yakovlev and A. P. Bilibin. The volume contains articles with data on the spectrum of action of biomyxin, its therapeutic forms, experimental therapy, clinical application, and effect on animal growth.

Prof L. Yakobson and associates report that they established the high activity of biomyxin in regard to various groups of microbes. It must be regretted, however, that the spectrum of action of biomyxin described in the article does not include the causative agents of the more dangerous infections.